

# Annexin V-PE Reagent

**CATALOG #:** 1014-200 200 assays (0.2 ml)  
1014-1000 1000 assays (1 ml)

**STORAGE CONDITIONS:** Store at 4°C. Do not freeze.

**SHELF LIFE:** 1 year under proper storage conditions

## DESCRIPTION:

Annexin V-PE is a bright fluorescent reagent for detecting of the early stages of apoptosis. During apoptosis, phosphatidylserine (PS) is translocated from the cytoplasmic face of the plasma membrane to the cell surface. Annexin V has a strong, Ca<sup>2+</sup>-dependent affinity for PS and therefore serves as a probe for detecting apoptosis. The one-step staining procedure takes only 10 minutes. In addition, the assay can be directly performed on live cells. Annexin V-PE stained cells can be detected by flow cytometry or fluorescence microscopy.

## ASSAY PROTOCOL:

### A. Incubation of cells with Annexin V-PE:

1. Induce apoptosis by desired methods.
2. Collect 1 x 10<sup>5</sup> cells by centrifugation.
3. Resuspend cells in 500 µl of 1X Annexin V Binding Buffer (Cat.#1035-100).
4. Add 1 µl of Annexin V-PE.
5. Incubate at room temperature for 5 min in the dark.  
Proceed to B or C below depending on method of analysis.

### B. Quantification by Flow Cytometry:

Analyze Annexin V-PE binding by flow cytometry (Ex = 488 nm; Em = 578 nm) using the phycoerythrin emission signal detector (usually FL2).

For analyzing adherent cells, gently trypsinize and wash cells once with serum-containing media before incubation with Annexin V-PE (A.3-5).

### C. Detection by Fluorescence Microscopy

1. Place the cell suspension from Step A.5 on a glass slide. Cover the cells with a glass coverslip.

For analyzing adherent cells, grow cells directly on a coverslip. Following incubation (A.5), invert coverslip on glass slide and visualize cells. The cells can also be washed and fixed in 2% formaldehyde before visualization. (Cells must be incubated with Annexin V-PE before fixation since any cell membrane disruption can cause nonspecific binding of Annexin V to PS on the inner surface of the cell membrane.)

2. Observe the cells under a fluorescence microscope using a rhodamine filter.

Cells which have bound Annexin V-PE will show bright orange-red staining in the plasma membrane.

## RELATED PRODUCTS:

### Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

### Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

### Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

### Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assay Kits

### Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

### Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

### Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β-Galactosidase Staining Kit & Luciferase Reporter Assay Kit

### Growth Factors and Cytokines

### Monoclonal and Polyclonal Antibodies