

Caspase-9 Colorimetric Substrate, LEHD-pNA

CATALOG #:	1076-200	200 assays (2 x 0.5 ml)
	1076-1000	1000 assays (5 x 1 ml)
LOT #:	_____	
STORAGE:	Store at -20°C, protected from light. Stable for 6 months	
MOL. WEIGHT:	675	
SEQUENCE:	Ac-Leu-Glu-His-Asp-pNA	
PURITY:	>98% by HPLC analysis.	

DESCRIPTION:

Ready-to-use colorimetric substrate for caspase-9/Mch6 and related caspases that recognize the amino acid sequence LEHD. Caspase-9 and related caspase activity can be quantified by spectrophotometric detection of free pNA ($\lambda = 400$ nm) after cleavage from the peptide substrate LEHD-pNA, using a spectrophotometer or multi-well plate reader.

ASSAY PROCEDURE:

1. Induce apoptosis in cells by desired method. Concurrently incubate a control culture *without* induction.
2. Count cells and pellet $1-5 \times 10^6$ cells.
3. Resuspend cells in 50 μ l of chilled Cell Lysis Buffer (Cat.# 1067-100, -400) and incubate cells on ice for 10 minutes.
4. Centrifuge for 1 min in a microcentrifuge (10,000 x g).
5. Transfer supernatant (cytosolic extract) to a fresh tube and put on ice.
6. Assay protein concentration.
7. Dilute 50-200 μ g protein to 50 μ l Cell Lysis Buffer for each assay.
8. Add 50 μ l of 2X Reaction Buffer (Cat.# 1068-20, -80) containing 10 mM DTT (Cat.# 1201-1) to each sample.
9. Add 5 μ l of the 4 mM substrate LEHD-pNA (200 μ M final conc.) into each tube and incubate at 37°C for 1-2 hour.
10. Read samples at 400 or 405-nm in a microtiter plate reader, or spectrophotometer using a 100- μ l micro quartz cuvet (Sigma), or dilute sample to 1 ml with Dilution Buffer (Cat.# 1066-100, -500) and using regular cuvet (note: Dilution of the samples proportionally decreases the reading).

You may also perform the entire assay directly in a 96-well plate.

Fold-increase in caspase activity can be determined by comparing these results with the level of the uninduced control.

Note: Background reading from cell lysates and buffers should be subtracted from the readings of both induced and the uninduced samples before calculating fold increase in caspase activity.

RELATED PRODUCTS:

Apoptosis Detection Kits & Reagents

- Annexin V Kits & Bulk Reagents
- Caspase Assay Kits & Reagents
- Mitochondrial Apoptosis Kits & Reagents
- Nuclear Apoptosis Kits & Reagents
- Apoptosis Inducers and Set
- Apoptosis siRNA Vectors

Cell Fractionation System

- Mitochondria/Cytosol Fractionation Kit
- Nuclear/Cytosol Fractionation Kit
- Membrane Protein Extraction Kit
- Cytosol/Particulate Rapid Separation Kit
- Mammalian Cell Extraction Kit
- FractionPREP Fractionation System

Cell Proliferation & Senescence

- Quick Cell Proliferation Assay Kit
- Senescence Detection Kit
- High Throughput Apoptosis/Cell Viability Assay Kits
- LDH-Cytotoxicity Assay Kit
- Bioluminescence Cytotoxicity Assay Kit
- Live/Dead Cell Staining Kit

Cell Damage & Repair

- HDAC Fluorometric & Colorimetric Assays & Drug Discovery Kits
- HAT Colorimetric Assay Kit & Reagents
- DNA Damage Quantification Kit
- Glutathione & Nitric Oxide Fluorometric & Colorimetric Assay Kits

Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β -Galactosidase Staining Kit & Luciferase Reporter Assay Kit

Growth Factors and Cytokines

Monoclonal and Polyclonal Antibodies