

## Recombinant Mouse Activin A

|                         |   |        |
|-------------------------|---|--------|
| <b>CATALOG #:</b>       | 4725-10   | 10 µg  |
|                         | 4725-100  | 100 µg |
|                         | 4725-1000   | 1 mg   |
| <b>ALTERNATE NAMES:</b> | Inhibin beta-1, FRP, FSH-releasing protein                                  |        |
| <b>SOURCE:</b>          | <i>E. coli</i>  |        |
| <b>PURITY:</b>          | ≥95% by HPLC, Reducing and Non-reducing SDS-PAGE, UV spectroscopy at 280 nm |        |
| <b>ENDOTOXIN LEVEL:</b> | ≤ 1 EU/µg protein   |        |
| <b>MOL. WEIGHT:</b>     | 26.2 kDa  |        |
| <b>FORMULATION:</b>     | Recombinant mouse Activin A is lyophilized with 0.02% TFA.                  |        |

### RECONSTITUTION:

Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water, which can then be further diluted to other aqueous solutions.

### STORAGE CONDITIONS:

Lyophilized product is very stable at -20°C. Reconstituted material should be aliquoted and frozen at -20°C. It is recommended that a carrier protein (0.1% HSA or BSA) is added for long term storage.

### DESCRIPTION:

Activin A is a member of the TGFbeta family of proteins produced by many cell types throughout development. Activins interact with Type I and Type II serine/threonine kinases to signal to SMAD proteins to regulate a variety of functions, including cell proliferation, differentiation, wound healing, apoptosis, metabolism, etc. Activin A is a homodimer of two beta A chains and is not biologically active until the N terminal propeptide is cleaved from each. Mouse Activin A has 100% amino acid sequence identity to human, rat, porcine, bovine and feline proteins. Recombinant mouse Activin A is a nonglycosylated homodimer, containing two 117 amino acid chains, with a total molecular weight of 26.2 kDa

### BIOLOGICAL ACTIVITY:

The activity is determined by the ability to induce cytotoxicity of MPC-11 cells and is typically less than 2 ng/mL.

### AA Sequence:

MGLECDGKVN ICCKKQFFVS FKDIGWNDWI IAPSGYHANY CEGECP SHIA  
GTSGSSLSFH STVINHYRMR GHSPFANLKS CCVPTKLRPM SMLYYDDGQN  
IHKKDIQNM IVEECGCS

### RELATED PRODUCTS:

- Activin A, human recombinant (Cat. No. 4724-10, -100, -1000)
- Activin A, rat recombinant (Cat. No. 4726-10, -100, -1000)

**FOR RESEARCH USE ONLY! Not to be used on humans.**