


## Anti-HMGA2 Antibody

<b>CATALOG NO:</b>	A1638-100 100 µl
<b>ALTERNATIVE NAMES:</b>	HMGA 2; BABL; High mobility group (nonhistone chromosomal) protein isoform I C; High mobility group (nonhistone chromosomal) protein isoform IC; High Mobility Group AT hook 2; High Mobility Group AT hook protein 2; High mobility group AT-hook protein 2; High mobility group protein HMGI C; High mobility group protein HMGI-C; High mobility group protein HMGIC; High mobility group protein I, isoform C; HMGA2; HMGA2_HUMAN; HMGI C; HMGIC; LIPO; Non histone chromosomal architectural protein HMGI C; Non histone chromosomal architectural protein HMGIC; Pygmy; STQTL9
<b>CONCENTRATION:</b>	1 mg/ml
<b>IMMUNOGEN:</b>	Peptide sequence around aa. 6~10(E-G-A-G-Q) derived from Human HMGA2.
<b>HOST/ISOTYPE:</b>	Rabbit IgG
<b>CLONALITY:</b>	Polyclonal
<b>MOL WEIGHT:</b>	18 kDa
<b>SPECIES REACTIVITY:</b>	Human, Mouse
<b>PURIFICATION:</b>	Affinity purification
<b>FORM:</b>	Liquid
<b>FORMULATION:</b>	Supplied in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02% sodium azide and 50% glycerol
<b>STORAGE CONDITIONS:</b>	For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles
<b>DESCRIPTION:</b>	Functions as a transcriptional regulator. Functions in cell cycle regulation through CCNA2.
<b>APPLICATION:</b>	WB 1:500-1:1000 ELISA (peptide) 1:20000-1:40000

**Note:** This information is only intended as a guide. The optimal dilutions must be determined by the user.



Western blot analysis of HeLa and 3T3 whole cell lysates using HMGA2 antibody

### RELATED PRODUCTS:

- GFAP Antibody (Cat. No. 3206)
- Vimentin Antibody (Cat. No. 3634)
- Fibronectin Antibody (Cat. No. 3630)
- FGF-2 Antibody (Cat. No. 5039)
- Anti-CD133 Antibody (Cat. No. A1622)
- Anti-CHRD1 Antibody (Cat. No. A1623)
- Anti-HES3 Antibody (Cat. No. A1636)

**FOR RESEARCH USE ONLY! Not to be used on humans.**